

Combining Ability Estimation for Fibre Quality Parameters in *Desi* Cotton (*Gossypium arboreum*)

Anil Kumar¹, H. V. Kalpande¹, V. N. Chinchane¹, Kuldeep Singh Chandrawat¹, Sunayana²
and P. B. Singh³

¹Department of Agricultural Botany, Vasantrao Naik Marathwada Krishi Vidyapeeth, Parbhani,
Maharashtra, India, 431-402

²Department of Genetics and Plant Breeding, CCSHAU Hisar, Haryana, India, 125004

³Department of Plant Breeding and Genetics, RCA (MPUAT), Udaipur, India, 313001
E-mail:- agusaiwal@gmail.com

Abstract

Present study was conducted to estimate combining ability and type of gene action for fibre quality parameters in *desi* cotton genotypes. Ten genotypes (six lines and four testers) and twenty four crosses were evaluated in randomized block design with three replications during kharif 2012-13. The results revealed that predominance of non-additive gene action was obtained for almost all the characters under study, such as ginning outturn, 2.5% length, fibre fineness, fibre strength uniformity ratio and short fibre index. The mean squares due to males were non-significant for all characters studied. The female lines exhibited significant differences for fibre strength only. Mean square due to crosses was significant for all the six characters indicating the importance of the studied traits. The estimates of GCA effects revealed that among lines, PA-08 was highest performing line for more number of characters followed by PA-720 and PA-255 while, among testers GAM-162 was the highest performing for more number of characters. These lines and testers being good general combiners can be used as donor parent for desirable genes regarding the mentioned traits.

Introduction

Most of the breeding programmes in cotton are designed for increasing yield and improve fibre quality because it is important for textile industry. The term general combining ability (GCA) is used to designate the average performance of a line in hybrid combinations and specific combining ability (SCA) as deviation in performance of a cross combination from that predicted on the basis of the general combining abilities of the parents involved in the cross^[1]. Combining ability analysis is used to compare the performance of breeding material^[5], allows estimation of different genetic components^[11] and reflects the performance of

lines, testers and crosses. Thus, the present investigation was conducted to identify superior parents and crosses in relation to fibre quality through L x T analysis.

Material and Methods

The present investigation was undertaken to study combining ability in *desi* cotton (*Gossypium arboreum* L.) for fibre quality traits in line x tester programme involving twenty four hybrid combinations which were derived by crossing six *arboreum* lines (PA-720, PA-08, PA-528, PA-532, PA-255 and PA-402) with four *arboreum* testers (AKA-7, GAM-162, Dwd-*arb*-10-1 and JLA-802). The

crosses and parents were evaluated at Cotton Research Station, Mahboob Baugh farm, Vasantao Naik Marathwada Krishi Vidyapeeth, Parbhani during *kharif* season of 2012-13.

The experiment was conducted in randomized block design in three replications. Data were recorded on five randomly selected plants per entry for all the six characters *viz.*, ginning out-turn (%), 2.5% span length (mm), micronaire/ fibre fineness (10^{-6} g/inch), fibre strength (g/tex) uniformity ratio (%) and short fibre index. All recorded datas were subjected to analysis of variance for testing the significance of treatments. Combining ability analysis and

the testing of significance of different genotypes was based on the standard^[9] procedure.

Results and Discussion

Analysis of variance of combining ability with respect to all six characters under study is summarized in Table-1. The mean squares due to males were non-significant for all characters studied. The female lines exhibited significant differences for fibre strength only. Mean square due to crosses was significant for all the six characters indicating the importance of the studied traits. Significant variance was observed for all the studied characters (excluding uniformity ratio) for line × tester interactions.

Table 1 Analysis of variance for combining Ability for different characters including parents

| Source | d.f. | Ginning outturn (%) | 2.5% SL length (%) | Fibre fineness/ Micronaire (μ g/inch) | Fibre strength (g/tex) | Uniformity ratio (%) | Short fibre index (%) |
|--------------|------|---------------------|--------------------|--|------------------------|----------------------|-----------------------|
| Replications | 2 | 3.453 | 0.017 | 0.008 | 0.557 | 0.007 | 0.050 |
| Crosses | 23 | 6.506** | 6.725** | 0.167** | 3.642** | 3.603* | 12.467** |
| Females | 5 | 6.567 | 11.078 | 0.170 | 9.398* | 7.125 | 5.325 |
| Males | 3 | 6.474 | 4.485 | 0.144 | 0.615 | 2.458 | 8.321 |
| M x F | 15 | 6.493** | 5.722** | 0.170** | 2.328** | 2.658 | 15.678** |
| Error | 66 | 0.561 | 0.699 | 0.051 | 0.984 | 1.961 | 0.326 |

By evaluation of GCA variances and SCA variances, it was observed that SCA variance was higher than GCA variances in all traits which indicate predominance of non-additive gene action^[10]. All the studied characters might be under non-additive gene

action (performed significance only for sca variances) except 2.5% span length and fibre strength (performed significance for both sca and gca variances) which might be due to both additive and non additive gene action (Table-2).

Table 2 Variances for General and Specific Combining Ability and Percent contribution of lines, testers and LxT for quality parameters in cotton

| Sr. No. | Character | δ^2 GCA | δ^2 SCA | δ^2 GCA / δ^2 SCA | Percent contributions of | | |
|---------|---|----------------|----------------|---------------------------------|--------------------------|---------|-------|
| | | | | | Lines | Testers | LxT |
| 1. | Ginning outturn (%) | 0.3973 | 1.9770** | 0.2009 | 21.94 | 12.97 | 65.07 |
| 2. | 2.5% SL length (%) | 0.4722* | 1.6743** | 0.2820 | 35.81 | 8.69 | 55.49 |
| 3. | Fibre fineness (Micronaire) (μ g/inch) | 0.0071 | 0.0396** | 0.1786 | 22.19 | 11.29 | 66.50 |
| 4. | Fibre strength (g/tex) | 0.2681* | 0.4478* | 0.5988 | 56.10 | 2.20 | 41.69 |
| 5. | Uniformity ratio (%) | 0.0107 | 0.0589** | 0.1817 | 42.98 | 8.89 | 48.11 |
| 6. | Short fibre index (%) | 0.4331 | 5.1170** | 0.0846 | 9.28 | 8.70 | 82.00 |

The predominance of additive gene action for fibre strength has been reported earlier^[2,4]. The ratio of δ^2 GCA / δ^2 SCA was less than unity for all the studied characters indicating preponderance of non additive gene action (dominance and epistasis). The contribution of line x tester was higher than both lines and testers for most of the characters except for fibre strength in which, contribution of line was more (56.10%). In comparison of both lines and testers it was found that contribution of lines to the total variability was more than testers for all the studied characters. Non-additive gene action for fibre quality traits viz., fibre length, fibre strength and micronaire value have been reported^[3, 6]. These results suggested that heterosis breeding was suitable for all the characters including fibre properties.

The estimates of general combining ability effects of females and males are presented in Table-3. Among lines, PA-720 (1.02) and PA-528 (0.60) while, among testers

GMA-162 (0.73) showed positive and significant gca effect for ginning outturn. The lines PA-08 (1.20) and PA-720 (0.80) while, among male parents, Dwd-arb-10-1 (0.64) displayed significant gca effect in positive direction respect of 2.5 % span length. The line PA-255 (0.18) exhibited highly positive significant gca effects and none of the testers exhibited significant gca effects for fibre fineness. As far as fibre strength is concerned, the line PA-08 (1.21) displayed significant GCA effects in positive direction while, none of the male parents exhibited significant gca effects. The line PA-08 (0.87) exhibited highly positive significant gca effects while, none of the testers exhibited significant gca effects for uniformity. Among female parents, PA-532 (-0.82) and PA-08 (-0.47) showed significant negative gca effects and among male parents, Dwd-arb-10-1 (-0.84) showed negative significant gca effect for short fibre index.

Table 3 Estimates of General Combining Ability (GCA) for Lines and Testers

| Parents | Ginning outturn (%) | 2.5% SL length (%) | Fibre fineness (Micronaire) (µg/inch) | Fibre strength (g/tex) | Uniformity ratio (%) | Short fibre index (%) |
|--------------|---------------------|--------------------|---------------------------------------|------------------------|----------------------|-----------------------|
| Lines | | | | | | |
| PA-720 | 1.02** | 0.80** | 0.01 | -0.85** | 0.62 | -0.32 |
| PA-08 | -0.90** | 1.20** | -0.09 | 1.21** | 0.87* | -0.47** |
| PA-528 | 0.60** | -0.21 | -0.14* | 0.54 | 0.37 | 0.17 |
| PA-532 | -0.05 | 0.30 | 0.08 | 0.49 | -0.62 | -0.82** |
| PA-255 | -0.71** | -0.79 | 0.18** | -0.38 | -0.12 | 0.97** |
| PA-402 | 0.04 | -1.31 | -0.04 | -1.00** | -1.12** | 0.47** |
| S.E.(Gi) | 0.216 | 0.241 | 0.6054 | 0.2864 | 0.404 | 0.1650 |
| S.E.(Gi-Gj) | 0.306 | 0.341 | 0.0925 | 0.4051 | 0.571 | 0.2333 |
| CD @5% | 0.435 | 0.485 | 0.1317 | 0.5766 | 0.813 | 0.3321 |
| CD @1% | 0.581 | 0.648 | 0.1758 | 0.7697 | 1.086 | 0.4433 |
| Testers | | | | | | |
| AKA-7 | -0.09 | 0.14 | 0.09 | -0.12 | 0.37 | 0.60** |
| GMA-162 | 0.73** | -0.45* | -0.00 | -0.19 | 0.20 | 0.49** |
| Dwd-arb-10-1 | -0.72** | 0.64** | 0.02 | 0.15 | -0.45 | -0.84** |
| JLA-802 | 0.08 | -0.32 | -0.11 | 0.15 | -0.12 | -0.25 |
| S.E.(Gi) | 0.176 | 0.197 | 0.0534 | 0.2339 | 0.330 | 0.1347 |
| S.E.(Gi-Gj) | 0.249 | 0.278 | 0.0755 | 0.3308 | 0.466 | 0.1905 |
| CD @5% | 0.355 | 0.396 | 0.1075 | 0.4708 | 0.664 | 0.2712 |
| CD @1% | 0.474 | 0.529 | 0.1435 | 0.6284 | 0.887 | 0.3620 |

The estimates of GCA effects revealed that among lines, PA-08 was highest performing line for more number of characters followed by PA-720 and PA-255 while, among testers GAM-162 was the highest performing for more number of characters. These lines and testers being good general combiners can be used as donor parent for desirable genes regarding the mentioned traits^[1].

The estimates of specific combining ability effects of combinations for various traits are given in Table 4. The hybrid PA-532 x JLA-802 (2.32) showed maximum significant positive SCA effects followed by cross PA-528 x AKA-7 (1.90), PA-402 x JLA-802 (1.53) and PA-402 x GAM-162 (1.23) were highest performing crosses for ginning outturn. The cross PA-720 x JLA-802 (2.32) showed maximum significant positive SCA effects followed by PA-255 x AKA-7 (1.55), PA-255 x

Dwd-arb-10-1 (1.28) and PA-528 x GAM-162 (1.15) for 2.5% span length. For Fibre fineness, the crosses viz., PA-532 x AKA-7 (-0.47) and PA-720 x JLA-802 (-0.33) showed significant negative SCA effect while, the crosses PA-720 x JLA-802 (1.52), PA-402 x AKA-7 (1.42) and PA-402 x JLA-802 (1.19) showed significant positive SCA effect for fibre strength. None of the crosses exhibited significant SCA effects for uniformity ratio. However, PA-402 x AKA-7 (1.29) and PA-08 x GAM-162 (1.20) were found to be highest performing crosses in positive direction. For short fibre index, the cross PA-720 x JLA-802 (-3.56) and PA-255 x AKA-7 (-3.44) recorded maximum significant negative SCA effects. Similar results related to GCA and SCA were also reported by earlier researchers^[3, 4, 8]. Anandan (2010), Bolek *et al.*, (2010), Karademr and Gencer (2010), Dewdar (2011) and DaiGang *et al.*, (2012) for these characters.

Table 4 Estimates of specific combining ability (SCA) for yield contributing characters

| Hybrids | Ginning outturn (%) | 2.5% SL length (%) | Fibre fineness (Micronaire) (µg/inch) | Fibre strength (g/tex) | Uniformity ratio (%) | Short fibre index (%) |
|-----------------------|---------------------|--------------------|---------------------------------------|------------------------|----------------------|-----------------------|
| PA-720 x JLA-802 | -1.458** | 2.358** | -0.332* | 1.525* | -0.625 | -3.558** |
| PA-528 x AKA-7 | 1.899** | -2.442** | 0.168 | -1.208* | -0.458 | 4.258** |
| PA-08 x GAM-162 | -0.816 | -0.342 | 0.079 | 0.242 | 1.208 | 0.292 |
| PA-532 x Dwd-arb-10-1 | 0.375 | 0.425 | 0.085 | -0.558 | -0.125 | -0.992** |
| PA-255 x JLA 802 | -0.649 | -0.342 | -0.224 | 0.250 | -0.875 | 0.292 |
| PA-402 x AKA-7 | 0.501 | -0.742 | 0.076 | 1.417* | 1.292 | -0.292 |
| PA-720 x GAM 162 | -0.470 | 0.258 | 0.154 | -1.233* | -0.042 | -0.158 |
| PA-528 x Dwd-arb-10-1 | 0.618 | 0.825 | -0.007 | -0.43 | -0.375 | 0.158 |
| PA-08 x JLA-802 | -1.344** | 0.783 | 0.226 | -0.075 | 0.625 | -1.458** |
| PA-532 x AKA -7 | -0.604 | 0.683 | -0.474** | -0.108 | -1.208 | -0.642 |
| PA-255 x GAM -162 | 0.091* | -0.817 | 0.004 | 0.342 | -0.542 | 0.19 |
| PA-402 x Dwd-arb-10-1 | 1.046* | -0.650 | 0.243 | -0.158 | 1.125 | 1.908** |
| PA-528 x JLA 802 | 1.014* | -1.342** | 0.101 | -0.525 | -0.375 | 1.442** |
| PA-720 x AKA-7 | 0.067 | 0.458 | 0.201 | 0.042 | -0.208 | 0.058 |
| PA-528 x GAM-162 | -0.211 | 1.158* | -0.221 | 0.392 | 0.458 | -1.608** |
| PA-08 x Dwd-arb-10-1 | -0.870 | -0.275 | -0.082 | 0.092 | 0.125 | 0.108 |
| PA-532 x JLA 802 | 2.322** | -0.842 | 0.301* | 0.050 | 1.125 | 0.042 |
| PA-255 x AKA-7 | -0.848 | 1.558** | -0.099 | 0.117 | 0.292 | -3.442** |
| PA-402 x GAM -162 | 1.230** | -1.542** | -0.121 | -0.033 | -0.042 | 2.592** |
| PA-720 x Dwd-arb-10-1 | -2.705** | 0.825 | -0.082 | -0.133 | -1.375 | 0.808* |
| PA-08 x AKA-7 | 0.115 | -0.617 | -0.074 | -1.225* | 0.125 | 3.242** |
| PA-532 x GAM-162 | -1.015* | 0.483 | 0.126 | -0.258 | 0.292 | 0.058 |
| PA-255 x Dwd-arb-10-1 | -0.634 | 1.283* | 0.104 | 0.292 | -1.042 | -1.308** |
| PA-402 x JLA-802 | 1.535** | -1.150* | -0.157 | 1.192* | 0.625 | -1.992** |
| S.E. ± | 0.4327 | 0.4828 | 0.1308 | 0.5729 | 0.8087 | 0.3300 |

References

1. Ahuja, S. L. and Dhayal, L. S. (2007). Combining ability estimates for yield and fibre quality traits in 4 x 13 line x tester crosses of *Gossypium hirsutum*. *Euphytica*, 153: 87-98.
2. Ali, I., Shakeel, A., Saeed, A., Nazeer, W., Zia, Z. U., Ahmad, S., Mahmood, K. and Malik, W. (2016). Combining ability analysis and heterotic studies for within-boll yield components and fibre quality in cotton. *The Journal of Animal & Plant Sciences*, 26(1): 156-162.
3. DaiGang Yang, Ma Xiong Feng, Zhou Xiao Jian, Zhang Xian Liang, Bai Feng Hu, Wang Hai Feng, Meng Qing Qin, Pei Xiao Yu and Yu Shuxun (2012). Correlation among combining ability, heterosis and genetic distance in upland cotton. [Chinese] *Cotton Science*, 24(3): 191-198.
4. Dewdar, M. D. H. (2011). Nature of combining abilities and genetic interpretation for some quantitative traits in Egyptian cotton. *Bulletin of Faculty of Agriculture*, Cairo University, 62(4): 418-424.
5. Griffing, B. (1956). Concept of general and specific combining ability in relation to diallel crossing system. *Australian Journal of Biological Science*, 9: 463-493.
6. Hassan, G., Mahood, G., Khan, N. U. and Razzaq, A. (1999). Combining ability and heterobeltiotic estimates in a diallel cross of cotton. *Sarhad Journal of Agriculture*, 15:563-568.
7. Hassan, G., Mahood, G., Razzaq, A. and Hayatullah (2000). Combining ability in inter-varietal crosses of Upland cotton (*Gossypium hirsutum* L.). *Sarhad Journal of Agriculture*, 16:407-410.
8. Karademr, E. and Gencer, O. (2010). Combining ability and heterosis for yield and fibre quality properties in cotton (*G. hirsutum* L.) obtained by half-diallel mating design. *Notulae Botanicae, Horti Agro botanici*, Cluj-Napoca., 38(1): 222-227.
9. Kempthorne, O. (1957). *An Introduction to Genetic Statistics*, New York, John Wiley and Sons, 1st Edn., pp. 456-471.
10. Tang, F. and W. Xiao (2013). Genetic effects and heterosis of within-boll yield components in upland cotton (*Gossypium hirsutum* L.). *Euphytica* 194(1): 41-51.
11. Verhalen, L. M. and Murray J. C. (1967). A diallel analysis of several fibre traits in upland cotton (*Gossypium hirsutum* L.). *Crop Science*, 7: 501-505.